arXiv:1601.06398v1 [math.AT] 24 Jan 2016

TWO-COMPLETE STABLE MOTIVIC STEMS OVER FINITE FIELDS

GLEN M. WILSON AND PAUL ARNE ØSTVÆR

Abstract. Let ℓ be a prime, and $q = p^n$ where p is a prime different from ℓ and 2. We show that the ℓ-completion of the nth stable homotopy group of spheres is a summand of the ℓ-completion of the $(n, 0)$ stable homotopy group of motivic spheres over the finite field with q elements $\mathbb{F}_q$. With this, and assisted by computer calculations, we are able to compute explicitly the two-complete stable motivic stems $\pi_{n,0}(\mathbb{F}_q)^\wedge_2$ for $0 \leq n \leq 20$ when $q \equiv 1 \bmod 4$ and for $0 \leq n \leq 18$ when $q \equiv 3 \bmod 4$.

1. Introduction

The homotopy groups of spheres belong to the most important and puzzling invariants in topology. See [Koc90] and the more recent works [Isa14b], [WX16] for amazing computer assisted ways of computing these invariants based on the Adams spectral sequence; this is a much studied approach to compute stable homotopy group of spheres [Ada95], [Rav86]. With two-primary coefficients, the second page of the Adams spectral sequence — converging to the two-completed stable homotopy groups of spheres — is comprised of Ext groups over the mod 2 Steenrod algebra,

$$E_2^{s,t} = \mathrm{Ext}^{s,t}_{\mathcal{A}^*}(\mathbb{F}_2, \mathbb{F}_2).$$

Extensive computer calculations of these Ext groups are carried out in [Bru93], [Bru16]. However, even if one knew completely the answer for the Ext groups in the Adams spectral sequence, one is still not finished with computing the stable homotopy groups of spheres. One needs to know in addition the differentials and all the group extensions hidden in the associated graded of the filtration. Only partial results have been obtained in spite of an enormous effort.

Given any field k, the stable motivic homotopy category $\mathcal{SH}(k)$ over k has the structure of a triangulated category, which encodes both algebro-geometric and topological information. An application of this setting is Voevodsky's proof of Milnor's conjecture on Galois cohomology [Voe03a]. Just as for the classical stable homotopy category $\mathcal{SH}$, it is an interesting and deep problem to compute the stable motivic homotopy groups of spheres $\pi_{p,q}(k)$ over k, i.e., $\mathcal{SH}(k)(\Sigma^{p,q}\mathbb{1}, \mathbb{1})$, where $\mathbb{1}$ denotes the motivic sphere spectrum over k. When k has finite mod 2 cohomological dimension, the motivic Adams spectral sequence, henceforth abbreviated to MASS, converges to the two-completed stable motivic stems

$$E_2^{f,(s,w)} = \mathrm{Ext}^{f,(s+f,w)}_{\mathcal{A}^{**}}(H^{**}, H^{**}) \Longrightarrow (\pi_{s,w}\mathbb{1})^\wedge_2.$$

Date: January 26, 2016.

2010 *Mathematics Subject Classification.* 14F42, 16-04, 18G15, 55T15.

Key words and phrases. Motivic Adams spectral sequence, stable motivic stems over finite fields, computer assisted motivic Ext group calculations.

This is a tri-graded spectral sequence, where $\mathcal{A}^{**}$ is the bi-graded mod 2 motivic Steenrod algebra, see e.g., [HKØ13] and [Voe03a], and H^{**} is the bi-graded mod 2 motivic cohomology ring of k. (We refer to Section 5 for the construction of MASS.) The calculational challenges consists basically of: (1) identify the motivic Ext groups, (2) determine the differentials, and (3) reconstruct the abutment from the filtration quotients.

Based on the MASS, Dugger and Isaksen have carried out calculations of the 2-complete stable motivic homotopy groups of spheres up to the 34-stem over the complex numbers [DI10]. Isaksen has extended this work largely up to the 70-stem [Isa14a, Isa14b]. We are led to wonder, how do the stable motivic homotopy groups vary for different base fields? Morel has given a complete description of the 0-line $\pi_{p,p}(k)$ in terms of Milnor-Witt K-theory [Mor12]. The 1-line $\pi_{p+1,p}(k)$ is determined by Hermitian and Milnor K-theory groups [RSØ16], generalizing the partial results obtained in [OØ14]. Ormsby has investigated the case of related invariants over p-adic fields [Orm11] and the rationals [OØ13], and Dugger and Isaksen have analyzed the case over the real numbers [DI15]. It is now possible to perform similar calculations over fields of positive characteristic, thanks to work on the motivic Steenrod algebra in positive characteristic [HKØ13]. In this paper we use computer assisted motivic Ext group calculations in tandem with theoretical arguments to determine stable motivic stems $\pi_{n,0}$ in weight zero over finite fields.

Let π_n^s denote the nth topological stable stem. Over the complex numbers, Levine showed there is an isomorphism $\pi_n^s \cong \pi_{n,0}(\mathbb{C})$ [Lev14, Cor. 2]. We obtain a similar result over the algebraic closure of a finite field, after ℓ-completion. For a prime ℓ and an Abelian group G, we write the ℓ-completion of G by $G^\wedge_\ell$.

Theorem A. Let p be a prime, and let $\overline{\mathbb{F}}_p$ denote the algebraic closure of the finite field with p elements. For any prime $\ell \neq p$ there is an isomorphism

$$\pi_{n,0}(\overline{\mathbb{F}}_p)^\wedge_\ell \cong (\pi_n^s)^\wedge_\ell.$$

Using theorem A, we are able to perform calculations in the MASS over finite fields at the prime 2 up to the 18-stem.

Theorem B. Let $\mathbb{F}_q$ be the finite field with q elements, where $q = p^i$ with p an odd prime. For any natural number $0 \leq n \leq 18$ there is an isomorphism

$$\pi_{n,0}(\mathbb{F}_q)^\wedge_2 \cong (\pi_n^s)^\wedge_2 \oplus (\pi_{n+1}^s)^\wedge_2.$$

In particular, the groups $\pi_{4,0}(\mathbb{F}_q)$ and $\pi_{12,0}(\mathbb{F}_q)$ are trivial.

In the case of a finite field $\mathbb{F}_q$ with $q \equiv 3 \bmod 4$, we require computer calculations to identify the structure of the E_2 page of the MASS. The details of these computer calculations, and a careful analysis of the differentials can be found in [KW15].

It is interesting to note that the pattern $\pi_{n,0}(\mathbb{F}_q)^\wedge_2 \cong (\pi_n^s)^\wedge_2 \oplus (\pi_{n+1}^s)^\wedge_2$ obtained in Theorem B does not hold in general. We show that if $q \equiv 5 \bmod 8$, then

$$\pi_{19,0}(\mathbb{F}_q)^\wedge_2 \cong (\mathbb{Z}/8 \oplus \mathbb{Z}/2) \oplus \mathbb{Z}/4 \text{ and } \pi_{20,0}(\mathbb{F}_q)^\wedge_2 \cong \mathbb{Z}/8 \oplus \mathbb{Z}/2.$$

We shall leave open for further investigations the question of whether or not an isomorphism $\pi_{n,0}(\mathbb{F}_q)^\wedge_2 \cong (\pi_n^s)^\wedge_2 \oplus (\pi_{n+1}^s)^\wedge_2$ holds when $q \equiv 3 \bmod 4$ and $n = 19, 20$.

2. The stable motivic homotopy category

We first sketch a construction of the stable motivic homotopy category that will be convenient for our purposes, and in the process, set our notation. Treatments of stable motivic homotopy theory can be found in [Ayo07, DRØ03, DLØ$^+$07, Jar00, Hu03].

2.1. **Base schemes.** A base scheme B is a Noetherian separated scheme of finite Krull dimension. We write Sm/B for the category of smooth schemes of finite type over B. Let $\mathrm{Pshv}(\mathrm{Sm}/B)$ denote the category of presheaves of sets on Sm/B. A space over B is a simplicial presheaf on Sm/B. The collection of spaces over B forms the category $\mathrm{Spc}(B) = \Delta^{op}\mathrm{Pshv}(\mathrm{Sm}/B)$, where morphisms are natural transformations of functors. We write $\mathrm{Spc}_*(B)$ for the category of pointed spaces. Note that $\mathrm{Spc}(B)$ is naturally equivalent to the category of presheaves on Sm/B with values in the category of simplicial sets $\underline{\mathrm{sSet}}$. We will occasionally switch between the two perspectives.

2.2. **The projective model structure.** The first model category structure we endow $\mathrm{Spc}(B)$ with is the projective model structure, see, e.g., [Bla01, Thm. 1.4], [DRØ03, Thm. 2.7], [Hir03, Thm. 11.6.1].

Definition 2.1. A map $f : X \to Y$ in $\mathrm{Spc}(B)$ is a (global) weak equivalence if for any $U \in \mathrm{Sm}/B$ the map $f(U) : X(U) \to Y(U)$ of simplicial sets is a weak equivalence. The projective fibrations are those maps $f : X \to Y$ for which $f(U) : X(U) \to Y(U)$ is a Kan fibration for any $U \in \mathrm{Sm}/B$. The projective cofibrations are those maps in $\mathrm{Spc}(B)$ which satisfy the left lifting property for trivial projective fibrations. This defines the projective model structure on $\mathrm{Spc}(B)$.

The category $\mathrm{Spc}(B)$ equipped with the projective structure is cellular, proper, and simplicial [Bla01, Thm. 1.4]. Furthermore, $\mathrm{Spc}(B)$ has the structure of a simplicial monoidal model category, with product $\times$ and internal hom $\underline{\mathrm{Hom}}$. We write $\mathrm{Map}(\mathcal{X}, \mathcal{Y})$ for the simplicial mapping space for spaces $\mathcal{X}$ and $\mathcal{Y}$.

Definition 2.2. For a smooth scheme X over B, we write rX for the representable presheaf of simplicial sets. For $U \in \mathrm{Sm}/B$, the simplicial set $rX(U)$ is given by $rX(U)_n = \mathrm{Sm}/B(U, X)$ for all $n \in \Delta$ where the face and degeneracy maps are the identity map. We will occasionally abuse notation and write X for rX.

The constant presheaf functor $c : \underline{\mathrm{sSet}} \to \mathrm{Spc}(B)$ associates to a simplicial set A the presheaf cA defined by $cA(U) = A$ for any $U \in \mathrm{Sm}/B$.

The functor c is a left Quillen functor when $\mathrm{Spc}(B)$ is equipped with the projective model structure. Its right adjoint $\mathrm{Ev}_B : \mathrm{Spc}(B) \to \underline{\mathrm{sSet}}$ satisfies $\mathrm{Ev}_B(X) = X(B)$. One can show that representable presheaves and constant presheaves in $\mathrm{Spc}(B)$ are cofibrant with the projective model structure.

2.3. **The Nisnevich local model structure.** Although representable presheaves embed into $\mathrm{Spc}(B)$, colimits which exist in Sm/B are not necessarily preserved in $\mathrm{Spc}(B)$. That is, if $X = \mathrm{colim}\, X_i$ in Sm/B, it need not be true that $rX = \mathrm{colim}\, rX_i$, e.g., $\mathrm{colim}(r\mathbb{A}^1 \leftarrow r\mathbb{G}_m \to r\mathbb{A}^1) \neq r\mathbb{P}^1$ (as one can check by applying the Picard group functor). To fix this, one introduces the Nisnevich topology on Sm/B.

Definition 2.3 (Nisnevich topology)**.** Let B be a base scheme. For any $X \in \mathrm{Sm}/B$, let $\mathcal{U} = \{U_i \to X\}$ be a finite family of étale maps in Sm/B. We say $\mathcal{U}$ is a Nisnevich

covering of X if for any $x \in X$ there exists a map $U_i \to X \in \mathcal{U}$ and a point $u \in U_i$ for which the induced map of residue fields $k(x) \to k(u)$ is an isomorphism. The Nisnevich covers determine a Grothendieck topology on Sm/B.

Definition 2.4 (Elementary distinguished squares)**.** An elementary distinguished square is a diagram in Sm/B

$$\begin{array}{ccc} V' & \longrightarrow & X' \\ \downarrow & & \downarrow f \\ V & \xrightarrow{j} & X \end{array}$$

for which f is an étale map, j is an open embedding, and $f^{-1}(X - V) \to X - V$ is an isomorphism, where these subschemes are given the reduced structure.

Proposition 2.5. The Nisnevich topology is the smallest topology containing the covers $\{V \to X, X' \to X\}$ which come from an elementary distinguished square.

Definition 2.6. For a pointed space $\mathcal{X}$, and $n \geq 0$, let $\pi_n\mathcal{X}$ denote the Nisnevich sheafification of the presheaf $U \mapsto \pi_n(\mathcal{X}(U))$.

Let W_{Nis} denote the class of maps $f : \mathcal{X} \to \mathcal{Y}$ for which $f_* : \pi_n\mathcal{X} \to \pi_n\mathcal{Y}$ is an isomorphism of Nisnevich sheaves for all $n \geq 0$. The Nisnevich local model structure on $\mathrm{Spc}_*(B)$ is the left Bousfield localization of the projective model structure with respect to W_{Nis}.

Definition 2.7. Let $W_{\mathbb{A}^1}$ be the class of maps $\pi_X : (X \times \mathbb{A}^1)_+ \to X_+$ for $X \in \mathrm{Sm}/B$. The motivic model structure on $\mathrm{Spc}_*(B)$ is the left Bousfield localization of the projective model structure with respect to $W_{Nis} \cup W_{\mathbb{A}^1}$. We write $\mathrm{Spc}_*^{\mathbb{A}^1}(B)$ for the category of pointed spaces equipped with the motivic model structure. The pointed motivic homotopy category $\mathcal{H}_*^{\mathbb{A}^1}(B)$ is the homotopy category of $\mathrm{Spc}_*^{\mathbb{A}^1}(B)$. Compare with [HKØ13, p. 5]. For pointed spaces X and Y, we write $[X, Y]$ for the set of maps $\mathcal{H}_*^{\mathbb{A}^1}(B)(X, Y)$.

There are two circles in the category of pointed spaces: the constant simplicial presheaf S^1 pointed at its 0-simplex and the representable presheaf $\mathbb{G}_m = \mathbb{A}^1 \setminus \{0\}$ pointed at 1. These determine a bi-graded family of spheres $S^{p,q} = (S^1)^{\wedge p-q} \wedge \mathbb{G}_m^{\wedge q}$.

Definition 2.8. For a pointed space X over B and natural numbers i and j, let $\pi_{i,j}X$ denote the set of maps $[S^{i,j}, X]$.

The category of pointed spaces $\mathrm{Spc}_*(B)$ equipped with the induced motivic model category structure has many good properties which make it amenable to Bousfield localization. In particular, $\mathrm{Spc}_*(B)$ is closed symmetric monoidal, pointed simplicial, left proper, and cellular.

2.4. **The stable Nisnevich local model structure.** With the unstable motivic model category in hand, we now construct the stable motivic model category using the general framework laid out in [Hov01].

Let T be a cofibrant replacement of $\mathbb{A}^1/\mathbb{A}^1 - \{0\}$. One can show that T is weak equivalent to $S^{2,1}$ in $\mathrm{Spc}_*^{\mathbb{A}^1}(B)$. The functor $T \wedge -$ on $\mathrm{Spc}_*^{\mathbb{A}^1}(B)$ is a left Quillen functor, which we may invert by creating a category of T-spectra.

Definition 2.9. A T-spectrum is a sequence of spaces $X_n \in \mathrm{Spc}_*^{\mathbb{A}^1}(B)$ equipped with structure maps $\sigma_n : T \wedge X_n \to X_{n+1}$. A map of T-spectra $f : X \to Y$ is a

collection of maps $f_n : X_n \to Y_n$ which is compatible with the structure maps. We write $\mathrm{Spt}_T(B)$ for the category of T-spectra of spaces.

The level model structure on $\mathrm{Spt}_T(B)$ is given by declaring a map $f : X \to Y$ to be a cofibration (resp. weak equivalence, fibration) if for every map $f_n : X_n \to Y_n$ is a cofibration (resp. weak equivalence, fibration) in the motivic model structure on $\mathrm{Spc}_*(B)$.

Definition 2.10. Let X be a T-spectrum. For integers i and j, the (i, j) stable homotopy sheaf of X is the sheaf $\pi_{i,j}X = \mathrm{colim}_n \pi_{i+2n,j+n}X_n$. A map $f : X \to Y$ is a stable weak equivalence if for all integers i and j, the induced maps $f_* : \pi_{i,j}X \to \pi_{i,j}Y$ are isomorphisms.

Definition 2.11. The stable model structure on $\mathrm{Spt}_T(B)$ is the model category where the weak equivalences are the stable weak equivalences and the cofibrations are the cofibrations in the level model structure. The fibrations are those maps with the right lifting property with respect to trivial cofibrations. We write $\mathcal{SH}(B)$ for the homotopy category of $\mathrm{Spt}_T(B)$ equipped with the stable model structure.

The stable model structure on $\mathrm{Spt}_T(B)$ can be realized as a left Bousfield localization of the level-wise model structure [Hov01, Def. 3.3].

Just as for the category Spt_{S^1} of simplicial S^1-spectra, there isn't a symmetric monoidal category structure on $\mathrm{Spt}_T(B)$ which lifts the smash product $\wedge$ in $\mathcal{SH}(B)$. One remedy is to use a category of symmetric T-spectra $\mathrm{Spt}_T^\Sigma(B)$. The construction of this category is given in [Hov01, Def. 7.7], [Jar00]. By [Hov01, Thm. 9.1], there is a zig-zag of Quillen equivalences from $\mathrm{Spt}_T^\Sigma(B)$ to $\mathrm{Spt}_T(B)$, so $\mathcal{SH}(B)$ is equivalent to the homotopy category of $\mathrm{Spt}_T^\Sigma(B)$ as well. The category $\mathcal{SH}(B)$ therefore is a symmetric monoidal, triangulated category, where the shift functor is given by $[1] = S^{1,0} \wedge -$.

In addition to the category of T-spectra, we will find it convenient to work with the category of $(\mathbb{G}_m, S^1)$ bi-spectra, see [Jar00, DLØ+07].

Definition 2.12. An S^1-spectrum over the base scheme B is a sequence of spaces $X_n \in \mathrm{Spc}_*(B)$ equipped with structure maps $\sigma_n : S^1 \wedge X_n \to X_{n+1}$. A map of S^1-spectra $f : X \to Y$ is a sequence of maps $f_n : X_n \to Y_n$ which is compatible with the structure maps. The collection of S^1-spectra with compatible maps between them forms a category $\mathrm{Spt}_{S^1}(B)$. Similar to the construction in definition 2.9, the stable model structure on $\mathrm{Spt}_{S^1}(B)$ is the left Bousfield localization of the level model structure on $\mathrm{Spt}_{S^1}(B)$ with respect to the motivic model structure on $\mathrm{Spc}_*(B)$ at the stable equivalences, cf. [Hov01, Def. 3.3].

Definition 2.13. In the projective model structure on $\mathrm{Spc}_*(B)$, the space $\mathbb{G}_m$ pointed at 1 is not cofibrant. We abuse notation, and write $\mathbb{G}_m$ for a cofibrant replacement of $\mathbb{G}_m$. A $(\mathbb{G}_m, S^1)$ bi-spectrum over B is a $\mathbb{G}_m$-spectrum of S^1-spectra. More concretely, a $(\mathbb{G}_m, S^1)$ bi-spectrum E is a bi-graded family of spaces $E_{m,n}$ with bonding maps $\sigma_{m,n} : S^1 \wedge E_{m,n} \to E_{m,n+1}$ and $\gamma_{m,n} : \mathbb{G}_m \wedge E_{m,n} \to E_{m+1,n}$ which are compatible, meaning that the following diagram commutes.

$$\begin{array}{ccccc}
\mathbb{G}_m \wedge S^1 \wedge E_{m,n} & & \xrightarrow{\tau \wedge E_{m,n}} & & S^1 \wedge \mathbb{G}_m \wedge E_{m,n} \\
\downarrow{\scriptstyle \mathbb{G}_m \wedge \sigma} & & & & \downarrow{\scriptstyle S^1 \wedge \gamma} \\
\mathbb{G}_m \wedge E_{m,n+1} & \xrightarrow{\gamma} & E_{m+1,n+1} & \xleftarrow{\sigma} & S^1 \wedge E_{m+1,n}
\end{array}$$

Thinking of $\mathrm{Spt}_{\mathbb{G}_m,S^1}$ as the category of $\mathbb{G}_m$-spectra of S^1-spectra, we first equip $\mathrm{Spt}_{\mathbb{G}_m,S^1}$ with the level model category structure with respect to the stable model category structure on $\mathrm{Spt}_{S^1}(B)$. The stable model category structure on $\mathrm{Spt}_{\mathbb{G}_m,S^1}$ is the left Bousfield localization at the class of stable equivalences.

There are evidently left Quillen functors $\Sigma^\infty_{S^1} : \mathrm{Spc}_*(B) \to \mathrm{Spt}_{S^1}(B)$ and $\Sigma^\infty_{\mathbb{G}_m} : \mathrm{Spt}_{S^1}(B) \to \mathrm{Spt}_{\mathbb{G}_m,S^1}(B)$. Additionally, the category $\mathrm{Spt}_{\mathbb{G}_m,S^1}(B)$ equipped with the stable model structure is Quillen equivalent to $\mathrm{Spt}_T(B)$.

Definition 2.14. To any spectrum of simplicial sets $E \in \mathrm{Spt}_{S^1}$ we may associate the constant S^1-spectrum cE over B with value E. That is, cE is the sequence of spaces cE_n with the evident bonding maps. For a simplicial spectrum E, we also write cE for the $(\mathbb{G}_m, S^1)$ bi-spectrum $\Sigma^\infty_{\mathbb{G}_m} cE$. This defines a left Quillen functor $c : \mathrm{Spt}_{S^1} \to \mathrm{Spt}_{\mathbb{G}_m,S^1}(B)$ with right adjoint given by evaluation at B. Compare with [Lev14, Lemma 6.5].

2.5. **Base change of stable model categories.**

Definition 2.15. Let $f : C \to B$ be a map of base schemes. Pull-back along f determines a functor $f^{-1} : \mathrm{Sm}/B \to \mathrm{Sm}/C$, which induces Quillen adjunctions $(f^*, f_*) : \mathrm{Spc}^{\mathbb{A}^1}_*(B) \to \mathrm{Spc}^{\mathbb{A}^1}_*(C)$ and $(f^*, f_*) : \mathrm{Spt}_T(B) \to \mathrm{Spt}_T(C)$. See e.g., [Mor05, §5] for more details.

Let Q (resp. R) denote the cofibrant (resp. fibrant) replacement functor in $\mathrm{Spt}_T(B)$. The derived functors $\mathbb{L}f^*$ and $\mathbb{R}f_*$ are given by the formulas $\mathbb{L}f^* = f^*Q$ and $\mathbb{R}f_* = f_*R$.

3. Comparison to the classical stable homotopy category

We start this section by recalling [Lev14, Thm. 1].

Theorem 3.1. If $B = \mathrm{Spec}(\mathbb{C})$, the map $\mathbb{L}c : \mathcal{SH} \to \mathcal{SH}(B)$ is fully faithful.

Proposition 3.2. Let $f : C \to B$ be a map of base schemes. The following diagram of stable homotopy categories commutes.



Proof. The result follows by considering the following diagram.



The left and right triangles are evidently commutative. The upper triangle commutes by the computation

$$(f^* c_B E)(U) = \operatorname*{colim}_{U \to f^{-1}V} c_B E(V) = E,$$

where the colimit is over the comma category $(U \downarrow f^{-1})$. Hence the entire diagram is commutative, and the result follows. □

Proposition 3.3. Let B be a base scheme equipped with a map $\mathrm{Spec}(\mathbb{C}) \to B$. Then $\mathbb{L}c : \mathcal{SH} \to \mathcal{SH}(B)$ is faithful.

Proof. For symmetric spectra X and Y, the map $\mathbb{L}c : \mathcal{SH}(X,Y) \to \mathcal{SH}(\mathbb{C})(cX, cY)$ factors through $\mathcal{SH}(B)(cX, cY)$ by proposition 3.2. Hence $\mathbb{L}c : \mathcal{SH}(X,Y) \to \mathcal{SH}(B)(cX, cY)$ is injective by theorem 3.1. □

Corollary 3.4. Write $W(\overline{\mathbb{F}}_p)$ for the ring of Witt vectors of $\overline{\mathbb{F}}_p$ and K for the fraction field of $W(\overline{\mathbb{F}}_p)$ [Ser79, II §6]. Then $\mathbb{L}c : \pi^s_n \to \pi_{n,0}(W(\overline{\mathbb{F}}_p))$ is an injection, as we have maps $W(\overline{\mathbb{F}}_p) \to K \to \mathbb{C}$.

4. Motivic cohomology

Spitzweck has constructed a spectrum $H\mathbb{Z}$ in $\mathrm{Spt}^{\Sigma}_T(B)$ which represents Levine's motivic cohomology $H^i_{mot}(X, n)$ — see [Lev01] — when B is the Zariski spectrum of a Dedekind domain. Spitzweck establishes enough nice properties of $H\mathbb{Z}$ for us to construct the MASS over general base schemes, and establish comparisons between the MASS over a Hensel local ring and its residue field.

4.1. Integral motivic cohomology.

Definition 4.1. Over the base scheme $\mathrm{Spec}(\mathbb{Z})$, the spectrum $H\mathbb{Z}_{\mathrm{Spec}(\mathbb{Z})}$ is defined in [Spi13, Def. 4.27]. For a general base scheme B, we define $H\mathbb{Z}_B$ to be $f^* H\mathbb{Z}_{\mathrm{Spec}(\mathbb{Z})}$ where $f : B \to \mathrm{Spec}(\mathbb{Z})$ is the unique map.

Let $B = \mathrm{Spec}(D)$ for D a Dedekind domain. For $X \in \mathrm{Sm}/B$, there is a canonical isomorphism $\mathcal{SH}(B)(\Sigma^\infty X_+, \Sigma^{i,n} H\mathbb{Z}) \cong H^i_{mot}(X, n)$, where H^i_{mot} denotes Levine's motivic cohomology defined using Bloch's cycle complex [Spi13, Cor. 7.19]. The isomorphism is functorial with respect to maps in Sm/B. Additionally, if $i : \{x\} \to B$ is the inclusion of a closed point with residue field $k(x)$, there is a commutative diagram.

$$\begin{array}{ccc} \mathcal{SH}(B)(\Sigma^\infty X_+, \Sigma^{i,n} H\mathbb{Z}) & \xrightarrow{\cong} & H^i_{mot}(X, n) \\ \downarrow & & \downarrow \\ \mathcal{SH}(k(x))(\mathbb{L}i^*\Sigma^\infty X_+, \Sigma^{i,n} H\mathbb{Z}) & \xrightarrow{\cong} & H^i_{mot}(\mathbb{L}i^* X, n) \end{array}$$

If the residue field $k(x)$ has positive characteristic, there is a canonical isomorphism of ring spectra $\mathbb{L}i^* H\mathbb{Z}_B \cong H\mathbb{Z}_{k(x)}$ [Spi13, Thm. 9.16].

Proposition 4.2. If $f : C \to B$ is a smooth map of base schemes, then $\mathbb{L}f^* H\mathbb{Z}_B \cong H\mathbb{Z}_C$.

Proof. Since f is smooth, $\mathbb{L}f^* = f^*$, see [Mor05, p. 44]. It is straightforward to see that $f^* H\mathbb{Z}_B \cong H\mathbb{Z}_C$, so the result follows. □

4.2. **Motivic cohomology with coefficients $\mathbb{Z}/\ell$.** For a prime ℓ, write $H\mathbb{Z}/\ell$ for the cofiber of the map $\ell\cdot : H\mathbb{Z} \to H\mathbb{Z}$ in $\mathcal{SH}(B)$. The spectrum $H\mathbb{Z}/\ell$ represents motivic cohomology with $\mathbb{Z}/\ell$ coefficients. For a smooth scheme X over B, we write $H^{*,*}(X;\mathbb{Z}/\ell)$ for the motivic cohomology of X with $\mathbb{Z}/\ell$ coefficients. The mod ℓ Steenrod algebra over B is the bi-graded algebra $\mathcal{A}^{**}(B) = \mathcal{SH}(B)(H\mathbb{Z}/\ell, H\mathbb{Z}/\ell)$. The dual mod ℓ Steenrod algebra over B is the Hopf algebroid $\mathcal{A}_{**}(B) = \pi_{**}(H\mathbb{Z}/\ell \wedge H\mathbb{Z}/\ell)$. When B is the spectrum of a ring R, we write $\mathcal{A}^{**}(R)$ for the Steenrod algebra instead of $\mathcal{A}^{**}(\mathrm{Spec}(R))$. We use the same convention for the dual Steenrod algebra and motivic cohomology.

Let $B = \mathrm{Spec}(D)$ be the spectrum of a Dedekind domain D in which ℓ is invertible. Spitzweck shows in [Spi13, Thm. 11.24] that the dual Steenrod algebra with $\mathbb{Z}/\ell$ coefficients $\mathcal{A}_{**}(D)$ has the expected structure as a Hopf algebroid, as given in [HKØ13, Thm. 5.5].

Definition 4.3. Let D be a Dedekind domain, and let C denote the set of sequences $(\epsilon_0, r_1, \epsilon_1, r_2, \ldots)$ with $\epsilon_i \in \{0,1\}$, $r_i \geq 0$, and only finitely many non-zero terms. The elements $\tau_i \in \mathcal{A}_{2\ell^i-1,\ell^i-1}(D)$ and $\xi_i \in \mathcal{A}_{2\ell^r-2,\ell^r-1}(D)$ are constructed in [Spi13, Cor. 11.23]. For any sequence $I = (\epsilon_0, r_1, \epsilon_1, r_2, \ldots)$ in C, write $\omega(I)$ for the element $\tau_0^{\epsilon_0}\xi_1^{r_1}\cdots$ and $(p(I), q(I))$ for the bi-degree of the operation $\omega(I)$.

We record Spitzweck's calculation of the dual Steenrod algebra [Spi13, Thm. 11.24] and the Steenrod algebra [Spi13, Remark 11.25] in the following propositions.

Proposition 4.4. Let D be a Dedekind domain. As an $H\mathbb{Z}/\ell$-module, there is a weak equivalence $\bigvee_{I\in B} \Sigma^{p(I),q(I)} H\mathbb{Z}/\ell \to H\mathbb{Z}/\ell \wedge H\mathbb{Z}/\ell$. The map is given by $\omega(I)$ on the factor $\Sigma^{p(I),q(I)} H\mathbb{Z}/\ell$.

Remark 4.5. Let D be a Dedekind domain in which ℓ is invertible, and consider the map $f : \mathbb{Z}[1/\ell] \to D$. A key observation in the proof of [Spi13, Thm. 11.24] is that the map $\mathbb{L}f^* : \mathcal{A}_{**}(\mathbb{Z}[1/\ell]) \to \mathcal{A}_{**}(D)$ satisfies $\mathbb{L}f^*\tau_i = \tau_i$ and $\mathbb{L}f^*\xi_i = \xi_i$ for all i. For a map $j : D \to \widetilde{D}$ of Dedekind domains in which ℓ is invertible, it follows that $\mathbb{L}j^*\tau_i = \tau_i$ and $\mathbb{L}j^*\xi_i = \xi_i$ for all i.

Proposition 4.6. Let D be a Dedekind domain in which ℓ is invertible. The Steenrod algebra over D is isomorphic to the dual of $\mathcal{A}_{**}(D)$, i.e., $\mathcal{A}^{**}(D) \cong \mathrm{Hom}(\mathcal{A}_{**}(D), H\mathbb{Z}/\ell_{**}(D))$.

5. Motivic Adams spectral sequence

The motivic Adams spectral sequence over a base scheme B may be defined using the appropriate notion of an Adams resolution, just as it is done classically [Ada95], [Rav86]. We recount the definition and establish some basic properties of the MASS under base change. We follow [DI10, §3] for the definition of the (cohomological) motivic Adams spectral sequence. See also [HKO11, §6].

Let $p \neq 2$ and ℓ be fixed primes, $p \neq \ell$. We will be interested in the specific case of the MASS over a field and over a discrete valuation ring (DVR) with residue field of characteristic p. We write H for the spectrum $H\mathbb{Z}/\ell$ over the base scheme B and $H^{**}(B)$ for the motivic cohomology of B with $\mathbb{Z}/\ell$ coefficients. The spectrum H is a ring spectrum and is cellular in the sense of [DI05] by [Spi13, Cor. 11.4].

5.1. **Construction of the mod ℓ MASS.**

Definition 5.1. Consider $X \in \mathrm{Spt}_T^\Sigma(B)$, and let $\overline{H}$ denote the spectrum in the cofibration sequence $\overline{H} \to \mathbb{1} \to H \to \Sigma\overline{H}$. The standard H-Adams resolution of X is the tower of cofibration sequences $X_{f+1} \to X_f \to W_f$ given by $X_f = \overline{H}^{\wedge f} \wedge X$ and $W_f = E \wedge X_f$. Compare with [Ada95, §15].



Definition 5.2. A collection of indices $\{(p_\alpha, q_\alpha) \in \mathbb{Z}^2 \mid \alpha \in A\}$ is said to be motivically finite if for any $\alpha \in A$ there are only finitely many $\beta \in A$ for which $p_\alpha \geq p_\beta$ and $2q_\alpha - p_\alpha \geq 2q_\beta - p_\beta$.

Definition 5.3 (H^{**}-MASS)**.** Let $X \in \mathrm{Spt}_T^\Sigma(B)$. An H^{**}-Adams resolution of X is a tower of cofibration sequences $X_{f+1} \to X_f \to W_f$ in $\mathcal{SH}(B)$ of the following form. Each spectrum W_f has a description $W_f \cong \vee_\alpha \Sigma^{p_\alpha, q_\alpha} H$ where the set of indices $\{(p_\alpha, q_\alpha)\}$ is motivically finite, and the induced map $H^{**}W_f \to H^{**}X_f$ is a surjection.



Proposition 5.4. Let $X \in \mathrm{Spt}_T^\Sigma(B)$ be cellular, and suppose $H^{**}(X)$ is a free H^{**}-module which is motivically finitely generated. The standard H-Adams resolution of X is an H^{**}-Adams resolution.

Proof. The conditions on X ensure that $H \wedge X$ is a motivically finite wedge of suspensions of H. Furthermore, the map $X_f \to H \wedge X_f = W_f$ induces a surjection $H^{**}W_f \to H^{**}X_f$. If $x : X_f \to H$ is a map in $H^{**}X_f$, the map $H \wedge x : H \wedge X_f \to H$ in $H^{**}W_f$ maps to x under j_f^*. □

Definition 5.5. Let $X \in \mathrm{Spt}_T^\Sigma(B)$ and let $\{X_f, W_f\}$ be the standard H-Adams resolution of X. The H-motivic Adams spectral sequence for X is obtained from the following exact couple.



The E_1 term of the MASS is $E_1^{f,(s,w)} = \pi_{s,w}W_f$. The index f is called the Adams filtration, s is the stem, and w is the motivic weight.

Definition 5.6. Let $X \in \mathrm{Spt}_T^\Sigma(B)$ and consider an Adams resolution $\{X_f, W_f\}$ of X. The Adams filtration of $\pi_{**}X$ is given by $F_i\pi_{**}X = \mathrm{im}(\pi_{**}X_i \to \pi_{**}X)$.

Proposition 5.7. Let $\mathfrak{S}$ denote the category of spectral sequences in the category of Abelian groups. The associated spectral sequence to the standard H-Adams

resolution defines a functor $MASS : \mathcal{SH}(B) \to \mathfrak{S}$. Furthermore, the motivic Adams spectral sequence is natural with respect to base change.

Proof. The construction of the standard H-Adams resolution is functorial. Given $X \to X'$ we get induced maps of H-Adams resolutions $\{X_f, W_f\} \to \{X'_f, W'_f\}$. As $\pi_{**}(-)$ is a triangulated functor, we get an induced map of the associated exact couples, and hence of spectral sequences $MASS(X) \to MASS(X')$.

Let $f : C \to B$ be a map of base schemes. The claim is that there is a natural transformation between $MASS : \mathcal{SH}(B) \to \mathfrak{S}$ and $MASS \circ \mathbb{L}f^* : \mathcal{SH}(B) \to \mathcal{SH}(C) \to \mathfrak{S}$. Let $X \in \mathcal{SH}(B)$ and let $\{X_f, W_f\}$ be the standard H_B-Adams resolution of X in $\mathcal{SH}(B)$. We may as well assume X is cofibrant, so $QX = X$ where Q is the cofibrant replacement functor. Let $\{X'_f, W'_f\}$ denote the standard H_C-resolution of $\mathbb{L}f^*X = f^*X$. Observe that $\{f^*X_f, f^*W_f\} = \{X'_f, W'_f\}$, since $f^*\mathbb{1} = \mathbb{1}$ and $f^*H_B = H_C$. We therefore have a map $\{\mathbb{L}f^*X_f, \mathbb{L}f^*W_f\} \to \{X'_f, W'_f\}$. Applying $\mathbb{L}f^* : \mathcal{SH}(B)(\mathbb{1}^{s,w}, -) \to \mathcal{SH}(C)(\mathbb{1}^{s,w}, \mathbb{L}f^*-)$ to $\{X_f, W_f\}$ gives a map of exact couples, and therefore a map $\Phi_X : MASS_B(X) \to MASS_C(\mathbb{L}f^*X)$. It is straightforward to verify that Φ determines a natural transformation. □

Corollary 5.8. For a map of base schemes $f : C \to B$, there is a map of motivic Adams spectral sequences $\Phi : MASS_B(\mathbb{1}) \to MASS_C(\mathbb{1})$. The map Φ is furthermore compatible with the induced map $\pi_{**}(\mathbb{1}_B) \to \pi_{**}(\mathbb{1}_C)$.

Proposition 5.9. Suppose $X \in \mathcal{SH}(B)$ satisfies the hypotheses of proposition 5.4. The MASS for X has E_2-page given by

$$E_2^{f,(s,w)} = \mathrm{Ext}^{f,(s+f,w)}_{\mathcal{A}^{**}(B)}(H^{**}X, H^{**}B)$$

with differentials $d_r : E_r^{f,(s,w)} \to E_r^{f+r,(s-1,w)}$ for $r \geq 2$.

Proof. The same argument given in the topological situation [Ada58, Thm. 2.1] goes through with minor modifications. □

Definition 5.10. Consider a base scheme $B = \mathrm{Spec}(D)$ for a Dedekind domain D. Let $\mathrm{Ext}(D)$ denote the tri-graded group $\mathrm{Ext}_{\mathcal{A}^{**}(B)}(H^{**}(B), H^{**}(B))$.

Corollary 5.11. If X and X' are in $\mathcal{SH}(B)$ which satisfy the hypotheses of proposition 5.4, and $X \to X'$ induces an isomorphism $H^{**}X' \to H^{**}X$, then the induced map $MASS(X) \to MASS(X')$ is an isomorphism of spectral sequences from the E_2-page onward.

Corollary 5.12. Let $f : C \to B$ be a map of base schemes. Suppose $\mathbb{L}f^* : H^{**}(B) \to H^{**}(C)$, $\mathbb{L}f^* : \mathcal{A}^{**}(B) \to \mathcal{A}^{**}(C)$, and $\mathbb{L}f^* : H^{**}X \to H^{**}(\mathbb{L}f^*X)$ are all isomorphisms. Then $MASS_B(X) \to MASS_C(\mathbb{L}f^*X)$ is an isomorphism of spectral sequences from the E_2 page onward.

5.2. **Convergence of MASS.**

Proposition 5.13. Let $B = \mathrm{Spec}(k)$ be the Zariski spectrum of a field k with characteristic $p \neq \ell$. For $X \in \mathcal{SH}(B)$, the motivic Adams spectral sequence converges $E_2^{f,(s,w)} \Rightarrow \pi_{s,w}X^\wedge_\ell$ under the following conditions: k has finite mod ℓ cohomological dimension, X is cellular and of finite type in the sense of [HKO11, §2], i.e., $H^{**}X$ has a set of generators which is motivically finite and $H^{**}X$ is free over $H^{**}(B)$.

Proof. The argument given in [HKO11] carries over to the positive characteristic case from the work of [HKØ13]. See [OØ14, Thm. 3.1] for the analogous argument for the motivic Adams-Novikov spectral sequence. □

Remark 5.14. The connectivity theorem of Morel is needed to identify what the MASS converges to. It was pointed out by Asok, Hoyois, Drew at the 2015 USC summer school on K-theory that the proof of the connectivity theorem for perfect fields has a gap. (More precisely, the published proof of [Mor04, Lem. 3.3.9] requires characteristic zero). Our calculations for finite fields are conditional upon a valid proof of Morel's connectivity theorem for finite fields. However, in [Mor12, Lemma 1.15], it is claimed that a version of Gabber's presentation lemma holds for finite fields, and that the connectivity theorem therefore holds. Should the reader be unsettled by this, we will show that our calculations over $\mathbb{F}_q$ at least compute $\pi_{**}(\widetilde{\mathbb{F}}_q)^\wedge_2$ where $\widetilde{\mathbb{F}}_q$ is the union of all field extensions $\mathbb{F}_{q^i}$ over $\mathbb{F}_q$ with i odd.

6. The stable stems $\pi_{n,0}$ over $\overline{\mathbb{F}}_p$

Let $W(\overline{\mathbb{F}}_p)$ denote the ring of Witt vectors of $\overline{\mathbb{F}}_p$, and let K denote the field of fractions of $W(\overline{\mathbb{F}}_p)$. Consider the maps $\overline{\mathbb{F}}_p \xleftarrow{\pi} W(\overline{\mathbb{F}}_p) \xrightarrow{j} \overline{K}$. The previous sections have set us up with enough machinery to compare the motivic Adams spectral sequences at a prime $\ell \neq p$ over the associated base schemes $\mathrm{Spec}(\overline{\mathbb{F}}_p)$, $\mathrm{Spec}(W(\overline{\mathbb{F}}_p))$, and $\mathrm{Spec}(\overline{K})$. We will often write the ring instead of the Zariski spectrum of the ring in our notation.

Proposition 6.1. The map $\pi : W(\overline{\mathbb{F}}_p) \to \overline{\mathbb{F}}_p$ induces an isomorphism of rings $\mathbb{L}\pi^* : H^{**}(W(\overline{\mathbb{F}}_p)) \to H^{**}(\overline{\mathbb{F}}_p)$, and $\mathbb{L}\pi^* : \mathcal{A}^{**}(W(\overline{\mathbb{F}}_p)) \to \mathcal{A}^{**}(\overline{\mathbb{F}}_p)$.

Proof. The spectrum H represents Levine's motivic cohomology with $\mathbb{Z}/\ell$ coefficients naturally, and base change induces an isomorphism $H^i(W(\overline{\mathbb{F}}_p), \mathbb{Z}/\ell(n)) \to H^i(\overline{\mathbb{F}}_p, \mathbb{Z}/\ell(n))$ by the rigidity theorem [Gei04, Thm. 1.2(3)].

From the description of $H \wedge H$ as an H-module in proposition 4.4 over $W(\overline{\mathbb{F}}_p)$, we see that the map $\mathcal{A}_{**}(W(\overline{\mathbb{F}}_p)) \to \mathcal{A}_{**}(\overline{\mathbb{F}}_p)$ will be an isomorphism if $\mathbb{L}\pi^*\tau_i = \tau_i$ and $\mathbb{L}\pi^*\xi_i = \xi_i$ for all i. This follows from remark 4.5. Now as the Steenrod algebra is the dual of $\mathcal{A}_{**}$ over fields and Dedekind domains, it follows that $\mathcal{A}^{**}(W(\overline{\mathbb{F}}_p)) \xrightarrow{\cong} \mathcal{A}^{**}(\overline{\mathbb{F}}_p)$. □

It is easily checked that $W(\overline{\mathbb{F}}_p)$ has the cardinality of the continuum by the construction in [Ser79, II §6]. Hence the cardinality of $\overline{K}$ is also that of the continuum. The classification of algebraically closed fields yields an isomorphism $\iota : \overline{K} \to \mathbb{C}$.

Proposition 6.2. The map $j : W(\overline{\mathbb{F}}_p) \to \overline{K}$ induces isomorphisms of graded rings $\mathbb{L}j^* : H^{**}(W(\overline{\mathbb{F}}_p)) \to H^{**}(\overline{K})$ and $\mathbb{L}j^* : \mathcal{A}^{**}(W(\overline{\mathbb{F}}_p)) \to \mathcal{A}^{**}(\overline{K})$. Furthermore, the maps $\mathbb{L}(\iota \circ j)^* : H^{**}(W(\overline{\mathbb{F}}_p)) \to H^{**}(\mathbb{C})$ and $\mathbb{L}(\iota \circ j)^* : \mathcal{A}^{**}(W(\overline{\mathbb{F}}_p)) \to \mathcal{A}^{**}(\mathbb{C})$ are isomorphisms.

Proof. The localization sequence in motivic cohomology [Gei05, §1.2] for the inclusion $\mathrm{Spec}(\overline{\mathbb{F}}_p) \to \mathrm{Spec}(W(\overline{\mathbb{F}}_p))$ is as follows.

$$\cdots H^{i-2,n-1}(\overline{\mathbb{F}}_p) \to H^{i,n}(W(\overline{\mathbb{F}}_p)) \to H^{i,n}(K) \to H^{i-1,n-1}(\overline{\mathbb{F}}_p) \cdots$$

Now since $H^{i,n}(\overline{\mathbb{F}}_p) = 0$ for $i \neq 0$, it follows that $H^{i,n}(W(\overline{\mathbb{F}}_p)) \to H^{i,n}(K)$ is injective. Rigidity implies $H^{**}(W(\overline{\mathbb{F}}_p)) \cong H^{**}(\overline{\mathbb{F}}_p)$, and since $\overline{K}$ is algebraically closed we conclude that the map $H^{**}(W(\overline{\mathbb{F}}_p)) \to H^{**}(K) \to H^{**}(\overline{K})$ is an isomorphism.

The same argument as in proposition 6.1 establishes that $\mathbb{L}j^* : \mathcal{A}^{**}(W(\overline{\mathbb{F}}_p)) \to \mathcal{A}^{**}(\overline{K})$ is an isomorphism. The maps $\mathbb{L}\iota^*$ are isomorphisms by the rigidity of $\mathcal{SH}(k)^\ell$ for algebraically closed fields of characteristic 0 [RØ08], so the result follows. □

Corollary 6.3. The maps $W(\overline{\mathbb{F}}_p) \xrightarrow{j} \overline{K} \xrightarrow{\iota} \mathbb{C}$, induce isomorphisms of E_2 pages

$$MASS(\mathbb{1}_{W(\overline{\mathbb{F}}_p)}) \xrightarrow{\mathbb{L}j^*} MASS(\mathbb{1}_{\overline{K}}) \xrightarrow{\mathbb{L}\iota^*} MASS(\mathbb{1}_{\mathbb{C}}).$$

Furthermore, the map $\mathbb{L}\pi^* : MASS(\mathbb{1}_{W(\overline{\mathbb{F}}_p)}) \to MASS(\mathbb{1}_{\overline{\mathbb{F}}_p})$ is an isomorphism at the E_2 page. Hence there are isomorphisms

$$E_2^{*,(*,*)}(\overline{\mathbb{F}}_p) \cong \mathrm{Ext}^{*,(*,*)}(\overline{\mathbb{F}}_p) \cong \mathrm{Ext}^{*,(*,*)}(\mathbb{C}).$$

Proof. The result follows from propositions 6.1, 6.2, and corollary 5.12. □

Proposition 6.4. Let B be the Zariski spectrum of $\overline{\mathbb{F}}_p$, $W(\overline{\mathbb{F}}_p)$, or $\mathbb{C}$. Let $\{X_f, W_f\}$ be an Adams resolution for $\mathbb{1}$ in $\mathcal{SH}(B)$, and define $\hat{\mathbb{1}} = \mathbb{1}/\operatorname{holim} X_f$. The MASS for $\mathbb{1}$ converges to $\pi_{**}\hat{\mathbb{1}}$, filtered by the Adams filtration.

Proof. The Adams filtration of $\pi_{**}\hat{\mathbb{1}}$ is exhaustive and Hausdorff by construction. The vanishing line in the MASS guarantees that the filtration is complete, hence convergence follows. □

We now give the proof for theorem A. That is, if $\ell \neq p$ is a prime, we show that the map $\mathbb{L}c : (\pi_n^s)^\wedge_\ell \to (\pi_{n,0}\mathbb{1}_{\overline{\mathbb{F}}_p})^\wedge_\ell$ is an isomorphism.

Proof. Consider the maps $\overline{\mathbb{F}}_p \leftarrow W(\overline{\mathbb{F}}_p) \to \overline{K}$. The induced maps on the MASS are compatible with the maps of homotopy groups $\pi_{**}\hat{\mathbb{1}}_{\overline{\mathbb{F}}_p} \leftarrow \pi_{**}\hat{\mathbb{1}}_{W(\overline{\mathbb{F}}_p)} \to \pi_{**}\hat{\mathbb{1}}_{\overline{K}}$. By corollary 6.3, the maps $MASS(\mathbb{1}_{\overline{\mathbb{F}}_p}) \leftarrow MASS(\mathbb{1}_{W(\overline{\mathbb{F}}_p)}) \to MASS(\mathbb{1}_{\mathbb{C}})$ are isomorphisms at the E_2 page. We thus have isomorphisms $\pi_{**}\hat{\mathbb{1}}_{\overline{\mathbb{F}}_p} \cong \pi_{**}\hat{\mathbb{1}}_{W(\overline{\mathbb{F}}_p)} \cong \pi_{**}\hat{\mathbb{1}}_{\mathbb{C}}$. By proposition 5.13, we identify $\pi_{**}\hat{\mathbb{1}}_{\overline{\mathbb{F}}_p} \cong \pi_{**}\mathbb{1}^\wedge_\ell$ and $\pi_{**}\hat{\mathbb{1}}_{\mathbb{C}} \cong \pi_{**}\mathbb{1}^\wedge_\ell$. Hence we have the following diagram.



Now as $\mathbb{L}c : (\pi_n^s)^\wedge_\ell \to (\pi_{n,0}\mathbb{1}_{\mathbb{C}})^\wedge_\ell$ is an isomorphism, it follows that all maps in the above diagram are isomorphisms. □

7. MASS for finite fields $\mathbb{F}_q$

We now analyze the 2-complete stable stems $\hat{\pi}_{**}(\mathbb{F}_q) = (\pi_{**}\mathbb{1}_{\mathbb{F}_q})^\wedge_2$ when q and 2 are relatively prime. The results of the previous section allow us to identify the nth classical 2-complete stable stem $\hat{\pi}^s_n = (\pi^s_n)^\wedge_2$ as a summand of $\hat{\pi}_{n,0}(\mathbb{F}_q)$. Using this, we are able to analyze the MASS for $\mathbb{F}_q$ in a range. For the remainder of this section, let H denote the spectrum representing motivic cohomology with $\mathbb{Z}/2$ coefficients.

7.1. **The E_2-page of MASS over $\mathbb{F}_q$.** Recall that the motivic cohomology of a field F has the description $H^{**}(F) \cong K^M_*(F)/2[\tau]$ where $K^M_n(F)/2$ is in degree (n,n), and τ is in degree $(0,1)$. As $K^M_1(\mathbb{F}_q) = \mathbb{F}_q^\times$ and $K^M_n(\mathbb{F}_q) = 0$ for $n \geq 2$, we conclude $H^{**}(\mathbb{F}_q) \cong \mathbb{F}_2[\tau,u]/(u^2)$ where u is the nontrivial element of $H^{1,1}(\mathbb{F}_q) \cong \mathbb{F}_q^\times/2$.

Proposition 7.1. If $q \equiv 1 \bmod 4$, the E_2-page of the MASS for $\mathbb{F}_q$ is given by

$$E_2^{*,(*,*)} = \mathrm{Ext}^{*,(*,*)}(\mathbb{F}_q) = E_2^{*,(*,*)}(\overline{\mathbb{F}}_p) \otimes_{\mathbb{F}_2} \mathbb{F}_2[\tau,u]/(u^2).$$

In the above, τ and u are of degree $(0,-1)$ and $(-1,-1)$, respectively.

Proof. Observe that $\mathcal{A}^{**}(\mathbb{F}_q) \cong \mathcal{A}^{**}(\overline{\mathbb{F}}_p) \otimes \mathbb{F}_2[\tau,u]/(u^2)$ and $H^{**}(\mathbb{F}_q) \cong H^{**}(\overline{\mathbb{F}}_p) \otimes \mathbb{F}_2[\tau,u]/(u^2)$. Since $\mathbb{F}_2[\tau,u]/(u^2)$ is flat as a module over $\mathbb{F}_2[\tau]$, a free resolution $H^{**}(\overline{\mathbb{F}}_p) \leftarrow P^\bullet$ determines a free resolution $H^{**}(\mathbb{F}_q) \leftarrow P^\bullet \otimes \mathbb{F}_2[\tau,u]/(u^2)$. As the canonical map

$$\mathrm{Hom}_{\mathcal{A}^{**}(\overline{\mathbb{F}}_p)}(-, H^{**}(\overline{\mathbb{F}}_p)) \otimes \mathbb{F}_2[\tau,u]/(u^2) \to \mathrm{Hom}_{\mathcal{A}^{**}(\mathbb{F}_q)}(- \otimes \mathbb{F}_2[\tau,u]/(u^2), H^{**}(\mathbb{F}_q)).$$

is a natural isomorphism, we conclude $\mathrm{Ext}(\overline{\mathbb{F}}_p) \otimes \mathbb{F}_2[\tau,u]/(u^2) \cong \mathrm{Ext}(\mathbb{F}_q)$. □

By the previous proposition, the irreducible elements of $\mathrm{Ext}(\mathbb{C})$ are also irreducible elements of $\mathrm{Ext}(\mathbb{F}_q)$ when $q \equiv 1 \bmod 4$. The only additional irreducible element in $\mathrm{Ext}(\mathbb{F}_q)$ is the class u. The irreducible elements of $\mathrm{Ext}(\mathbb{F}_q)$ up to stem $s = 21$ can be found in table 1. These were obtained by consulting [Isa14b, Table 8], and independently verified by [KW15].

Elt.	Filtr. (f,s,w)	Elt.	Filtr. (f,s,w)	Elt.	Filtr. (f,s,w)
u	$(0,-1,-1)$	c_0	$(3,8,5)$	e_0	$(4,17,10)$
τ	$(0,0,-1)$	Ph_1	$(5,9,5)$	P^2h_1	$(9,17,9)$
h_0	$(1,0,0)$	Ph_2	$(5,11,6)$	f_0	$(4,18,10)$
h_1	$(1,1,1)$	d_0	$(4,14,8)$	P^2h_2	$(9,19,10)$
h_2	$(1,3,2)$	h_4	$(1,15,8)$	c_1	$(3,19,11)$
h_3	$(1,7,4)$	Pc_0	$(7,16,9)$	$[\tau g]$	$(4,20,11)$

TABLE 1. The irreducible elements of $\mathrm{Ext}(\mathbb{F}_q)$ with $q \equiv 1 \bmod 4$ in stem $s \leq 21$

If $q \equiv 3 \bmod 4$, the E_2 page of the MASS for $\mathbb{F}_q$ can be identified in a range using the ρ-Bockstein spectral sequence (ρ-BSS). Here $\rho = [-1]$ is the non-zero class in $H^{1,1}(\mathbb{F}_q) \cong \mathbb{F}_q^\times/2$, since -1 is not a square in $\mathbb{F}_q^\times$. We briefly describe the construction of the ρ-BSS, and refer the reader to [DI15], [OØ13], [Orm11] for more details.

Let $\mathcal{C}$ be the cobar construction corresponding to the Hopf algebroid

$$(\mathbb{F}_2[\tau,\rho]/(\rho^2), \mathcal{A}_{**}(\mathbb{F}_q)).$$

The filtration of $\mathcal{C}$ given by $0 \subseteq \rho\mathcal{C} \subseteq \mathcal{C}$ determines a spectral sequence, which in this case is just the long exact sequence associated to the short exact sequence of complexes

$$0 \to \rho\mathcal{C} \to \mathcal{C} \to \mathcal{C}/\rho\mathcal{C} \to 0.$$

Note that $\rho\mathcal{C}$ and $\mathcal{C}/\rho\mathcal{C}$ are both isomorphic to the cobar construction over $\mathbb{C}$. Hence we have a long exact sequence

$$\cdots \rho\,\mathrm{Ext}^{i,(*,*)}(\mathbb{C}) \longrightarrow \mathrm{Ext}^{i,(*,*)}(\mathbb{F}_q) \longrightarrow \mathrm{Ext}^{i,(*,*)}(\mathbb{C}) \xrightarrow{d_1} \rho\,\mathrm{Ext}^{i+1,(*,*)}(\mathbb{C}) \cdots$$

The differential d_1 satisfies the Leibniz rule, so it suffices to identify the differential on irreducible elements. We describe these differentials in the following proposition.

Proposition 7.2. In the ρ-BSS for $\mathbb{F}_q$ with $q \equiv 3 \bmod 4$, every irreducible element x of $\mathrm{Ext}(\mathbb{C})$ in stem $s \leq 19$ other than τ has $d_1 x = 0$. Also, $d_1\tau = \rho h_0$ and $d_1([\tau g]) = \rho h_2 e_0$. Here, $[\tau g]$ is the irreducible element of $\mathrm{Ext}(\mathbb{C})$ in stem 20, weight 11, and filtration 4.

Proof. The differential d_1 vanishes on the classes h_0, h_1, c_0, Ph_1, d_0, Pc_0, e_0, P^2h_1 since there is nothing for them to hit. The remaining differentials were identified by computer calculation in [KW15]. □

Example 7.3. Since $d_1 h_1 = 0$, we conclude $d_1\tau h_1 = \rho h_0 h_1 = 0$, as $h_0 h_1$ vanishes in $\mathrm{Ext}(\mathbb{C})$. Hence there is a class $[\tau h_1] \in \mathrm{Ext}^{1,(1,0)}(\mathbb{F}_q)$ which is irreducible.

With this analysis of the ρ-BSS for $\mathbb{F}_q$ with $q \equiv 3 \bmod 4$, one can determine the structure of $\mathrm{Ext}(\mathbb{F}_q)$ as a graded Abelian group up to stem 20. In [DI15], Dugger and Isaksen are additionally able to determine the product structure of $\mathrm{Ext}(\mathbb{R})$ from the ρ-BSS. Some of their techniques apply over $\mathbb{F}_q$. We instead use a computer calculation of $\mathrm{Ext}(\mathbb{F}_q)$ which includes the product structure in our analysis of the stable stems.

Proposition 7.4. The irreducible elements of $\mathrm{Ext}(\mathbb{F}_q)$ up to stem $s = 21$ are given in table 2.

Proof. This was obtained by computer calculation in [KW15]. □

7.2. **Differentials on the E_2 page.** We now use the unit map $\mathbb{1} \to H\mathbb{Z}$ to identify some differentials in $MASS(\mathbb{1}_{\mathbb{F}_q})$.

Proposition 7.5. The MASS for $H\mathbb{Z}$ over $\mathbb{F}_q$ when $q \equiv 1 \bmod 4$ has E_2 page given by

$$E_2^{*,(*,*)} \cong \mathbb{F}_2[\tau, u, h_0]/(u^2).$$

Let ν_2 denote the 2-adic valuation, and write $\epsilon(q)$ for $\nu_2(q-1)$. The differentials for $MASS(H\mathbb{Z})$ are given by

$$d_{\epsilon(q)+\nu_2(i)}\tau^i = u\tau^{i-1}h_0^{\epsilon(q)+\nu_2(i)}.$$

When $q \equiv 3 \bmod 4$, the E_2 page of $MASS(H\mathbb{Z})$ is given by

$$E_2^{*,(*,*)} \cong \mathbb{F}_2[\tau^2, \rho, h_0]/(\rho h_0, \rho^2) \oplus \rho\tau\mathbb{F}_2[\tau^2, \rho, h_0]/(\rho^2)$$

Elt.	Filtr. (f,s,w)	Elt.	Filtr. (f,s,w)	Elt.	Filtr. (f,s,w)
ρ	$(0,-1,-1)$	$[\tau c_0]$	$(3,8,4)$	$[\tau P c_0]$	$(7,16,8)$
$[\rho\tau]$	$(0,-1,-2)$	Ph_1	$(5,9,5)$	e_0	$(4,17,10)$
$[\tau^2]$	$(0,0,-2)$	$[\tau Ph_1]$	$(5,9,4)$	P^2h_1	$(9,17,9)$
h_0	$(1,0,0)$	Ph_2	$(5,11,6)$	$[\tau P^2h_1]$	$(9,17,8)$
h_1	$(1,1,1)$	$[\tau h_0h_3^2]$	$(3,14,7)$	f_0	$(4,18,10)$
$[\tau h_1]$	$(1,1,0)$	d_0	$(4,14,8)$	P^2h_2	$(9,19,10)$
h_2	$(1,3,2)$	$[\tau h_0^2d_0]$	$(6,14,7)$	c_1	$(3,19,11)$
$[\tau h_2^2]$	$(2,6,3)$	h_4	$(1,15,8)$	$[\tau c_1]$	$(3,19,10)$
h_3	$(1,7,4)$	$[\tau h_0^7h_4]$	$(8,15,7)$	$[\rho\tau g]$	$(4,19,10)$
$[\tau h_0^3h_3]$	$(4,7,3)$	Pc_0	$(7,16,9)$	$[\tau^2 g]$	$(4,20,10)$
c_0	$(3,8,5)$				

TABLE 2. The irreducible elements of $\mathrm{Ext}(\mathbb{F}_q)$ with $q \equiv 3 \bmod 4$ in stem $s \leq 21$

Let $\lambda(q) = \nu_2(q^2-1)$. For $q \equiv 3 \bmod 4$, the differentials for $MASS(H\mathbb{Z})$ are given by

$$d_{\lambda(q)+\nu_2(i)}\tau^{2i} = \rho\tau^{2i-1}h_0^{\lambda(q)+\nu_2(i)}.$$

Proof. Observe that the triangle $H\mathbb{Z} \xrightarrow{\times 2} H\mathbb{Z} \to H\mathbb{Z}/2$ gives rise to the following Adams resolution of $H\mathbb{Z}$.



We therefore may identify the E_1 page of $MASS(H\mathbb{Z})$ with $\oplus_{f\geq 0}\pi_{**}H\mathbb{Z}/2$. We can infer the differentials, as $MASS(H\mathbb{Z}) \Rightarrow (H\mathbb{Z}_{**})^\wedge_2$, and $(H\mathbb{Z}_{**})^\wedge_2$ is known by étale cohomology calculations, see e.g., [Sou79, III.1.4, IV.2]. Specifically we have

$$(H\mathbb{Z}_{i,j})^\wedge_2 \cong \begin{cases} \mathbb{Z}_2 & \text{if } i=j=0 \\ \mathbb{Z}/(q^j-1)^\wedge_2 & \text{if } i=-1, j\geq 1 \\ 0 & \text{otherwise.} \end{cases}$$

□

7.3. **Stable stems over $\mathbb{F}_q$.** We now begin an analysis of the differentials in the MASS to identify the two-complete stable stems over $\mathbb{F}_q$. To assist the reader with the computations presented below, figure 1 and figure 4 display E_2 page charts of the MASS over $\mathbb{F}_q$. Further charts can be found in [KW15].

Proposition 7.6. Let $q = p^n$ where p is a prime different from 2. The group $\hat{\pi}^s_n$ is a summand of $\hat{\pi}_{n,0}(\mathbb{F}_q)$ for $n \geq 0$.

Proof. The map $\mathbb{L}c : \pi^s_n \to \pi_{n,0}\mathbb{1}_{\overline{\mathbb{F}}_p}$ factors through $\pi_{n,0}\mathbb{1}_{\mathbb{F}_q}$. Passing to the 2-completion, theorem A implies the composition $\hat{\pi}^s_n \to \hat{\pi}_{n,0}(\mathbb{F}_q) \to \hat{\pi}_{n,0}(\overline{\mathbb{F}}_p)$ is an isomorphism. Hence the result. □

For small values of $n \geq 0$ we have $\hat{\pi}_{n,0}(\mathbb{F}_q) \cong \hat{\pi}^s_n \oplus \hat{\pi}^s_{n+1}$. This pattern fails when $n = 19$ and $q \equiv 5 \bmod 8$.

Proposition 7.7. When $q \equiv 5 \bmod 8$ and $0 \leq n \leq 18$, there is an isomorphism $\hat{\pi}_{n,0}(\mathbb{F}_q) \cong \hat{\pi}^s_n \oplus \hat{\pi}^s_{n+1}$.

Proof. The irreducible elements of $\mathrm{Ext}(\mathbb{F}_q)$ in this range are tabulated in proposition 7.4. All differentials d_r for $r \geq 2$ vanish on $h_0, h_1, h_3, c_0, Ph_1, d_0, Pc_0, P^2h_1$ since there is nothing for them to hit. As $\hat{\pi}_{3,0}(\mathbb{F}_q)$ must contain $\hat{\pi}^s_3 \cong \mathbb{Z}/8$ as a summand, we conclude $d_2(h_2) = 0$. As Ph_2 equals the Massey product $\langle h_3, h_0^4, h_2\rangle$ the differential $d_2(Ph_2) = 0$, since $d_2(h_2) = 0$. Similarly, as $P^2h_2 = \langle h_3, h_0^4, h_2\rangle$ we calculate $d_2P^2h_2 = 0$.

The differential $d_2(h_4) = h_0h_3^2$ is deduced from the comparison $MASS(\mathbb{F}_q) \to MASS(\overline{\mathbb{F}}_q) \cong MASS(\mathbb{C})$. This comparison also shows $d_3(h_0h_4)$ is non-zero. In order to have $\hat{\pi}^s_{14}$ as a summand of $\hat{\pi}_{14,0}$, we must have $d_3(h_0h_4) = h_0d_0$. Similarly, $d_2(e_0) = h_1^2d_0$, $d_2(f_0) = h_0^2e_0$. Note that $d_4(h_0^4h_4) = 0$ since there is nothing for it to hit. The elements in weight 0 are all of the form $\tau^i x$ or $u\tau^{i-1}x$ where x is not a multiple of τ and of weight i. The differentials of the elements in weight 0 are now readily identified by using the Leibniz rule. Since $\hat{\pi}^s_n$ is a summand of $\hat{\pi}_{n,0}(\mathbb{F}_q)$ for all n, we see that there are no hidden extensions for $n \leq 18$, so the result follows. □

Proposition 7.8. When $q \equiv 5 \bmod 8$, $\hat{\pi}_{19,0}(\mathbb{F}_q) \cong (\mathbb{Z}/8 \oplus \mathbb{Z}/2) \oplus \mathbb{Z}/4$ and $\hat{\pi}_{20,0}(\mathbb{F}_q) \cong \mathbb{Z}/8 \oplus \mathbb{Z}/2$.

Proof. We must have $d_2[\tau g] = 0$ for dimension reasons. Since $[\tau g]$ has weight 11, the class $\tau^{11}[\tau g]$ is in $E^{4,(20,0)}$. Using proposition 7.5, we calculate $d_2\tau^{11}[\tau g] = u\tau^{10}h_0^2[\tau g] \neq 0$. This resolves all of the differentials in the 19- and 20-stem, and the calculation of the 19-stem follows. As $\hat{\pi}^s_{20} \cong \mathbb{Z}/8$ must be a summand of $\hat{\pi}_{20,0}(\mathbb{F}_q)$, we conclude there is a hidden extension from $u\tau^{11}h_2^2h_4 = u\tau^{11}h_3^3$ to $\tau^{12}h_2e_0$. The calculation of the 20-stem now follows. □

Remark 7.9. Note that over $\mathbb{F}_q$ with $q \equiv 5 \bmod 8$, the map $\mathbb{L}c\{g\}$ is detected by $u\tau^{11}h_3^3$ which is in Adams filtration 3, and not 4. But over $\overline{\mathbb{F}}_q$, $\mathbb{L}c\{g\}$ is in Adams filtration 4.

Proposition 7.10. When $q \equiv 1 \bmod 8$ and $0 \leq n \leq 20$, there is an isomorphism $\hat{\pi}_{n,0}(\mathbb{F}_q) \cong \hat{\pi}^s_n \oplus \hat{\pi}^s_{n+1}$.

Proof. The same argument as in proposition 7.7 goes through in this case as well. The difference is that when $q \equiv 1 \bmod 8$, $d_2\tau = 0$ by proposition 7.5, and therefore $d_2\tau^{11}[\tau g] = 0$. This determines all of the differentials needed to compute up to the 20-stem. We note that there are no possible hidden extensions in the 19- or 20-stem in this case. The result then follows. □

Proposition 7.11. When $q \equiv 3 \bmod 4$ and $0 \leq n \leq 18$, there is an isomorphism $\hat{\pi}_{n,0}(\mathbb{F}_q) \cong \hat{\pi}^s_n \oplus \hat{\pi}^s_{n+1}$.

Proof. The differentials d_r for $r \geq 2$ vanish on the following generators for degree reasons: $[\rho\tau]$, ρ, h_0, h_1, h_3, $[\tau h_2^2]$, $[\tau c_0]$, $[\tau Ph_1]$, d_0, $[\tau Pc_0]$, $[\tau P^2h_1]$. Since $\hat{\pi}^s_1 \cong \mathbb{Z}/2$ is a summand of $\hat{\pi}_{1,0}(\mathbb{F}_q)$, we must have $d_r[\tau h_1] = 0$ for all $r \geq 2$. Since $\hat{\pi}^s_3 \cong \mathbb{Z}/8$ is a summand of $\hat{\pi}_{3,0}(\mathbb{F}_q)$, we must have $d_2(h_2) = 0$. Just as in proposition 7.7, we conclude $d_2(h_4) = h_0h_3^2$, $d_2(e_0) = h_1^2d_0$, and $d_2(f_0) = h_0^2e_0$ by comparison to $MASS(\overline{\mathbb{F}}_q)$. Also, we determine $d_r[\tau c_1] = 0$ for $r \geq 2$ by comparing with $MASS(\overline{\mathbb{F}}_q)$, as the class $[\tau c_1]$ must be a permanent cycle.

The one exceptional case is $d_3(h_0h_4)$. Here we must have $d_3(h_0h_4) = h_0d_0+\rho h_1d_0$ in order for $\hat{\pi}^s_{14} = \mathbb{Z}/2 \oplus \mathbb{Z}/2$ to be a summand of $\hat{\pi}_{14,0}(\mathbb{F}_q)$.

The elements in weight 0 are all of the form $([\tau^2])^i x$ or $[\rho\tau][\tau^2]^{i-1}x$ where x is not a multiple of τ^2 and weight $2i$, or of the form $\rho[\tau^2]^i x$ if x is not a multiple of τ^2 and of weight $2i+1$. The differentials of the elements in weight 0 are now readily identified by using the Leibniz rule. Since $\lambda(q) = \nu_2(q^2-1) \geq 3$, we have $d_2\tau^2 = 0$. This is sufficient to ensure that for elements x in stem $n \leq 18$ there are no non-trivial differentials of the form $d_r[\tau^2]^i x = \rho\tau^{2i-1}h_0^r x$ when $[\tau^2]^i x$ has weight 0. □

Remark 7.12. It is unclear whether $d_2[\tau^2 g] = [\rho\tau g]$ or $d_2[\tau^2 g] = 0$. This is all that obstructs the identification of the stems $\hat{\pi}_{19,0}(\mathbb{F}_q)$ and $\hat{\pi}_{20,0}(\mathbb{F}_q)$.

7.4. **Base change for finite fields.**

Proposition 7.13. Let $q = p^n$ where p is an odd prime. For a field extension $f : \mathbb{F}_q \to \mathbb{F}_{q^i}$ with i odd, the induced map $\mathbb{L}f^* : H^{**}(\mathbb{F}_q) \to H^{**}(\mathbb{F}_{q^i})$ and $\mathbb{L}f^* : \mathcal{A}^{**}(\mathbb{F}_q) \to \mathcal{A}^{**}(\mathbb{F}_{q^i})$ are isomorphisms.

Proof. The claim follows by checking on étale cohomology. The map on cohomology is determined by $H^1_{et}(\mathbb{F}_q; \mu_2) \to H^1_{et}(\mathbb{F}_{q^i}; \mu_2)$ which is just the induced map $\mathbb{F}_q^\times/2 \to \mathbb{F}_{q^i}^\times/2$. So long as i is odd, this map is an isomorphism. □

Corollary 7.14. For $q = p^n$ with p an odd prime, the induced map $MASS(\mathbb{F}_q) \to MASS(\mathbb{F}_{q^i})$ is an isomorphism of spectral sequences whenever i is odd.

Proposition 7.15. Let $q = p^n$ with p an odd prime. Let $\widetilde{\mathbb{F}}_q$ denote the union of the field extensions $\mathbb{F}_{q^i}$ over $\mathbb{F}_q$ with i odd. The field extension $f : \mathbb{F}_q \to \widetilde{\mathbb{F}}_q$ induces isomorphisms $\mathbb{L}f^* : H^{**}(\mathbb{F}_q) \to H^{**}(\widetilde{\mathbb{F}}_q)$ and $\mathbb{L}f^* : \mathcal{A}^{**}(\mathbb{F}_q) \to \mathcal{A}^{**}(\widetilde{\mathbb{F}}_q)$. Hence the map $MASS(\mathbb{F}_q) \to MASS(\widetilde{\mathbb{F}}_q)$ is an isomorphism of spectral sequences.

Proof. This follows by a colimit argument, using proposition 7.13. □

Corollary 7.16. Assuming Morel's connectivity theorem holds for $\mathbb{F}_q$, for any integers i and j, there is an isomorphism $\hat{\pi}_{i,j}(\mathbb{F}_q) \cong \hat{\pi}_{i,j}(\widetilde{\mathbb{F}}_q)$.

Proposition 7.17. Let $q = p^n$ with p an odd prime. For a field extension $f : \mathbb{F}_q \to \mathbb{F}_{q^i}$ with i even, the map $\mathbb{L}f^* : H^{1,*}(\mathbb{F}_q) \to H^{1,*}(\mathbb{F}_{q^i})$ is the trivial map.

Proof. The map is determined by $\mathbb{L}f^* : H^{1,1}(\mathbb{F}_q) \to H^{1,1}(\mathbb{F}_{q^i})$ which is just the map $\mathbb{F}_q^\times/2 \to \mathbb{F}_{q^i}^\times/2$. However, any non-square $x \in \mathbb{F}_q^\times$, will be a square in $\mathbb{F}_{q^i}^\times$ when i is even. □

Corollary 7.18. Let $q = p^n$ with p an odd prime. For a field extension $f : \mathbb{F}_q \to \mathbb{F}_{q^i}$ with i even, the induced map $MASS(\mathbb{F}_q) \to \mathbb{F}_{q^i}$ kills the class u, (respectively ρ and $[\rho\tau]$) and all of their multiples at the E_2 page.

Proof. The induced map of cobar complexes is determined from proposition 7.17, which shows the classes u (respectively ρ and $[\rho\tau]$) are killed under base change. □

8. Implementation of motivic Ext group calculations

The computer calculations used in this paper were done with the program available from [KW15] at `https://github.com/glenwilson/MassProg`. The program is written in python, and calculates $\mathrm{Ext}(F)$ for the fields $\mathbb{C}$, $\mathbb{R}$, and $\mathbb{F}_q$ by producing a minimal resolution of $H^{**}(F)$ by $\mathcal{A}^{**}(F)$ modules in a range. With this complex in hand, the program then produces its dual and calculates cohomology in each degree.

To calculate a free resolution of $H^{**}(F)$ of $\mathcal{A}^{**}(F)$-modules, we first need the program to efficiently perform calculations in $\mathcal{A}^{**}(F)$. The mod 2 motivic Steenrod algebra is generated by the squaring operations Sq^i and the cup products $\alpha \cup -$ for $\alpha \in H^{**}(F)$. These generators satisfy Adem relations, which are given in [HKØ13, Thm. 5.1], [Voe03b, Thm. 10.2]. Additionally, one needs the commutation relations $\mathrm{Sq}^{2i}\tau = \tau\,\mathrm{Sq}^{2i} + \tau\rho\,\mathrm{Sq}^{2i-1}$ for $i > 0$ and $\mathrm{Sq}^{2i+1}\tau = \tau\,\mathrm{Sq}^{2i+1} + \rho\,\mathrm{Sq}^{2i} + \rho^2\,\mathrm{Sq}^{2i-1}$ for $i \geq 0$ which are obtained from the Cartan formula. With these relations, the program can calculate the canonical form of any element of $\mathcal{A}^{**}$, that is, as a sum of monomials $\alpha \cdot \mathrm{Sq}^I$ where $\alpha \in H^{**}(F)$ and I is an admissible sequence.

With the algebra of $\mathcal{A}^{**}(F)$ available to the program, it then proceeds to calculate a minimal resolution of $H^{**}(F)$ by $\mathcal{A}^{**}(F)$ modules. This is where a great deal of computational effort is spent. To clarify what a minimal resolution is in practice, let $\prec$ denote the order on $\mathbb{Z}\times\mathbb{Z}$ given by $(m_1, n_1) \prec (m_2, n_2)$ if and only if $m_1+n_1 < m_2+n_2$, or $m_1+n_1 = m_2+n_2$ and $n_1 < n_2$. The reader is encouraged to compare this definition with the definition in [McC01, Def. 9.3], and consult [Bru09] for detailed calculations of a minimal resolution for the Adams spectral sequence of topology.

Definition 8.1. A resolution of $H^{**}(F)$ by $\mathcal{A}^{**}(F)$ modules $H^{**}(F) \leftarrow P^\bullet$ is a minimal resolution if the following conditions are satisfied.

1. Each module P^i has an ordered basis $h_i(j)$ such that if $j \leq k$ then $\deg h_i(j) \prec \deg h_i(k)$.
2. $\mathrm{im}(h_i(j)) \notin \mathrm{im}(\langle h_i(j) \,|\, j < k\rangle)$
3. $\deg h_i(j)$ is minimal with respect to degree in the order $\prec$ over all elements in $P^{i-1} \setminus \mathrm{im}(\langle h_i(j) \,|\, j < k\rangle)$.

The computer program calculates the first n maps and modules in a minimal resolution up to bi-degree $(2n, n)$. With this, it then calculates the dual of the resolution by applying the functor $\mathrm{Hom}_{\mathcal{A}^{**}(F)}(-, H^{**}(F))$ to the resolution $P^\bullet$. With the cochain complex $\mathrm{Hom}_{\mathcal{A}^{**}(F)}(P^\bullet, H^{**}(F))$ in hand, the program calculates cohomology in each degree, i.e., $\mathrm{Ext}^{f,(s+f,w)}(\mathbb{F}_q)$.

As the program calculates an explicit resolution of $H^{**}(F)$, the products of elements in $\mathrm{Ext}(F)$ can be obtained from the composition product, cf. [McC01, Thm. 9.5].

9. Charts

The weight 0 part of the E_2 page of the mod 2 MASS over $\mathbb{F}_q$ is depicted in figures 1, 4 according to the case $q \equiv 1 \bmod 4$ or $q \equiv 3 \bmod 4$. The weight 0 part of the E_∞ page of the mod 2 MASS over $\mathbb{F}_q$ can be found in figures 2, 3, 5.

In each chart, a circular or square dot in grading (s, f) represents a generator of the $\mathbb{F}_2$ vector space in the graded piece of the spectral sequence. The square dots

are used to indicate that the given element is divisible by u, ρ, or $\rho\tau$, depending on the case. Circular dots denote elements which are not divisible by u, ρ, or $\rho\tau$. In the chart 5, there is an oval dot which corresponds to the class with representative $\tau^8\rho h_1 d_0 \equiv \tau^8 h_0 d_0$, as the class $\rho h_1 d_0 + h_0 d_0$ is killed.

The labels in the chart correspond to those elements coming from irreducible elements of $\mathrm{Ext}(\mathbb{F}_q)$. However, most of these elements must be multiplied by the appropriate power of τ or $[\tau^2]$ to land in the weight 0 part of the spectral sequence. We leave this off in the notation, as the weights of these elements are listed in tables 1 and 2.

We indicate that the product of a given class by h_0 with a solid, vertical line. In the case $q \equiv 3 \bmod 4$, multiplication by ρh_1 plays an important role, so non-zero products by ρh_1 are indicated by dashed vertical lines. In particular, when $q \equiv 3 \bmod 4$, multiplication by 2 in $\hat{\pi}_{**}(\mathbb{F}_q)$ is detected by multiplication by $h_0+\rho h_1$. The lines of slope 1 indicate multiplication by τh_1 or $[\tau h_1]$ depending on the case.

Dotted lines are used in two separate instances in these charts. The first use is in figure 3, where dotted lines indicate hidden extensions by h_0 and τh_1. The other instance is in figure 5 to indicate an unknown d_2 differential.

Additional charts can be found in [KW15].



FIGURE 1. E_2 page of MASS for $\mathbb{F}_q$ with $q \equiv 1 \bmod 4$, weight 0



FIGURE 2. E_∞ page of MASS for $\mathbb{F}_q$ with $q \equiv 1 \bmod 8$, weight 0



FIGURE 3. E_∞ page of MASS for $\mathbb{F}_q$ with $q \equiv 5 \bmod 8$, weight 0



FIGURE 4. E_2 page of MASS for $\mathbb{F}_q$ with $q \equiv 3 \bmod 4$, weight 0



FIGURE 5. E_∞ page of MASS for $\mathbb{F}_q$ with $q \equiv 3 \bmod 4$, weight 0

Acknowledgments.

The work on this paper was initiated during the program "A Special Semester in Motivic Homotopy Theory" organized by Marc Levine in 2014; we gratefully acknowledge the hospitality and support from Universität Duisburg-Essen.

The first author would like to thank Charles Weibel, Dan Isaksen, and Kyle Ormsby for helpful conversations regarding this work. This paper constitutes a portion of the first author's PhD thesis. The second author was supported by the RCN project Special Geometries, no. 239015.

References

[Ada58] J. F. Adams, *On the structure and applications of the Steenrod algebra*, Comment. Math. Helv. **32** (1958), 180–214. MR 0096219 (20 #2711)

[Ada95] ———, *Stable homotopy and generalised homology*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1995, Reprint of the 1974 original. MR 1324104 (96a:55002)

[Ayo07] J. Ayoub, *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. II*, Astérisque (2007), no. 315, vi+364 pp. (2008). MR 2438151 (2009m:14007)

[Bla01] B. A. Blander, *Local projective model structures on simplicial presheaves*, K-Theory **24** (2001), no. 3, 283–301. MR 1876801 (2003c:18014)

[Bru93] R. R. Bruner, Ext *in the nineties*, Algebraic topology (Oaxtepec, 1991), Contemp. Math., vol. 146, Amer. Math. Soc., Providence, RI, 1993, pp. 71–90. MR 1224908 (94a:55011)

[Bru09] ———, *An adams spectral sequence primer*, math.wayne.edu/~rrb/papers/adams.pdf (2009).

[Bru16] ———, *The cohomology of the mod 2 steenrod algebra*, math.wayne.edu/~rrb/papers/cohom.pdf (2016).

[DI05] D. Dugger and D. C. Isaksen, *Motivic cell structures*, Algebr. Geom. Topol. **5** (2005), 615–652. MR 2153114 (2007c:55015)

[DI10] ———, *The motivic Adams spectral sequence*, Geom. Topol. **14** (2010), no. 2, 967–1014. MR 2629898 (2011e:55024)

[DI15] ———, *Low dimensional Milnor-Witt stems over* $\mathbb{R}$, arXiv:1502.01007 (2015).

[DLØ+07] B. I. Dundas, M. Levine, P. A. Østvær, O. Röndigs, and V. Voevodsky, *Motivic homotopy theory*, Universitext, Springer-Verlag, Berlin, 2007, Lectures from the Summer School held in Nordfjordeid, August 2002. MR 2334212 (2008k:14046)

[DRØ03] B. I. Dundas, O. Röndigs, and P. A. Østvær, *Motivic functors*, Doc. Math. **8** (2003), 489–525 (electronic). MR 2029171 (2004m:55011)

[Gei04] T. Geisser, *Motivic cohomology over Dedekind rings*, Math. Z. **248** (2004), no. 4, 773–794. MR 2103541 (2006c:14026)

[Gei05] ———, *Motivic cohomology, K-theory and topological cyclic homology*, Handbook of K-theory. Vol. 1, 2, Springer, Berlin, 2005, pp. 193–234. MR 2181824 (2006j:14025)

[Hir03] P. S. Hirschhorn, *Model categories and their localizations*, Mathematical Surveys and Monographs, vol. 99, American Mathematical Society, Providence, RI, 2003. MR 1944041 (2003j:18018)

[HKO11] P. Hu, I. Kriz, and K. M. Ormsby, *Remarks on motivic homotopy theory over algebraically closed fields*, J. K-Theory **7** (2011), no. 1, 55–89. MR 2774158 (2012b:14040)

[HKØ13] M. Hoyois, S. Kelly, and P. A. Østvær, *The motivic Steenrod algebra in positive characteristic*, arXiv:1305.5690, to appear in JEMS (2013).

[Hov01] M. Hovey, *Spectra and symmetric spectra in general model categories*, J. Pure Appl. Algebra **165** (2001), no. 1, 63–127. MR 1860878 (2002j:55006)

[Hu03] P. Hu, *S-modules in the category of schemes*, Mem. Amer. Math. Soc. **161** (2003), no. 767, viii+125. MR 1950209 (2004b:55016)

[Isa14a] D. C. Isaksen, *Classical and motivic Adams charts*, arXiv:1401.4983 (2014).

[Isa14b] ———, *Stable stems*, arXiv:1407.8418 (2014).

[Jar00] J. F. Jardine, *Motivic symmetric spectra*, Doc. Math. **5** (2000), 445–553 (electronic). MR 1787949 (2002b:55014)

[Koc90] S. O. Kochman, *Stable homotopy groups of spheres*, Lecture Notes in Mathematics, vol. 1423, Springer-Verlag, Berlin, 1990, A computer-assisted approach. MR 1052407 (91j:55016)

[KW15] Fu. K. and G. M. Wilson, *Cohomology of the Steenrod algebra over* $\mathbb{R}$, math.rugers.edu/~wilson47/ (2015).

[Lev01] M. Levine, *Techniques of localization in the theory of algebraic cycles*, J. Algebraic Geom. **10** (2001), no. 2, 299–363. MR 1811558 (2001m:14014)

[Lev14] ———, *A comparison of motivic and classical stable homotopy theories*, J. Topol. **7** (2014), no. 2, 327–362. MR 3217623

[McC01] J. McCleary, *A user's guide to spectral sequences*, Cambridge University Press, 2001.

[Mor04] F. Morel, *An introduction to* $\mathbb{A}^1$*-homotopy theory*, Contemporary developments in algebraic K-theory, ICTP Lect. Notes, XV, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004, pp. 357–441 (electronic). MR 2175638 (2006m:19007)

[Mor05] ———, *The stable* $\mathbb{A}^1$*-connectivity theorems*, K-Theory **35** (2005), no. 1-2, 1–68. MR 2240215 (2007d:14041)

[Mor12] ———, $\mathbb{A}^1$*-algebraic topology over a field*, Lecture Notes in Mathematics, vol. 2052, Springer, Heidelberg, 2012. MR 2934577

[OØ13] K. M. Ormsby and P. A. Østvær, *Motivic Brown-Peterson invariants of the rationals*, Geom. Topol. **17** (2013), no. 3, 1671–1706. MR 3073932

[OØ14] ———, *Stable motivic* π_1 *of low-dimensional fields*, Adv. Math. **265** (2014), 97–131. MR 3255457

[Orm11] K. M. Ormsby, *Motivic invariants of p-adic fields*, J. K-Theory **7** (2011), no. 3, 597–618. MR 2811717 (2012h:14057)

[Rav86] D. C. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, Pure and Applied Mathematics, vol. 121, Academic Press, Inc., Orlando, FL, 1986. MR 860042 (87j:55003)

[RØ08] O. Röndigs and P. A. Østvær, *Rigidity in motivic homotopy theory*, Math. Ann. **341** (2008), no. 3, 651–675. MR 2399164 (2009e:14034)

[RSØ16] O. Röndigs, M. Spitzweck, and P. A. Østvær, *The first stable homotopy groups of motivic spheres*, in preparation (2016).

[Ser79] J.-P. Serre, *Local fields*, Graduate Texts in Mathematics, vol. 67, Springer-Verlag, New York-Berlin, 1979, Translated from the French by Marvin Jay Greenberg. MR 554237 (82e:12016)

[Sou79] C. Soulé, *K-théorie des anneaux d'entiers de corps de nombres et cohomologie étale*, Invent. Math. **55** (1979), no. 3, 251–295. MR 553999 (81i:12016)

[Spi13] M. Spitzweck, *A commutative* $\mathbb{P}^1$*-spectrum representing motivic cohomology over Dedekind domains*, arXiv:1207.4078 (2013).

[Voe03a] V. Voevodsky, *Motivic cohomology with* $\mathbf{Z}/2$*-coefficients*, Publ. Math. Inst. Hautes Études Sci. (2003), no. 98, 59–104. MR 2031199 (2005b:14038b)

[Voe03b] ———, *Reduced power operations in motivic cohomology*, Publ. Math. Inst. Hautes Études Sci. (2003), no. 98, 1–57. MR 2031198 (2005b:14038a)

[WX16] G. Wang and Z. Xu, *The algebraic Atiyah-Hirzebruch spectral sequence of real projective spectra*, arXiv:1601.02185 (2016).

Department of Mathematics, Rutgers, The State University of New Jersey, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.

E-mail address: glen.m.wilson@rutgers.edu

Department of Mathematics, University of Oslo, Norway.

E-mail address: paularne@math.uio.no